thiopurine S-methyltransferase deficiency

Thiopurine S-methyltransferase (TPMT) deficiency is a condition characterized by significantly reduced activity of an enzyme that helps the body process drugs called thiopurines. These drugs, which include 6-thioguanine, 6-mercaptopurine, and azathioprine, inhibit (suppress) the body's immune system. Thiopurine drugs are used to treat some autoimmune disorders, including Crohn disease and rheumatoid arthritis, which occur when the immune system malfunctions. These drugs are also used to treat several forms of cancer, particularly cancers of blood-forming tissue (leukemias) and cancers of immune system cells (lymphomas). Additionally, thiopurine drugs are used in organ transplant recipients to help prevent the immune system from attacking the transplanted organ.

A potential complication of treatment with thiopurine drugs is damage to the bone marrow (hematopoietic toxicity). Although this complication can occur in anyone who takes these drugs, people with TPMT deficiency are at highest risk. Bone marrow normally makes several types of blood cells, including red blood cells, which carry oxygen; white blood cells, which help protect the body from infection; and platelets, which are involved in blood clotting. Damage to the bone marrow results in myelosuppression, a condition in which the bone marrow is unable to make enough of these cells. A shortage of red blood cells (anemia) can cause pale skin (pallor), weakness, shortness of breath, and extreme tiredness (fatigue). Low numbers of white blood cells (neutropenia) can lead to frequent and potentially life-threatening infections. A shortage of platelets (thrombocytopenia) can cause easy bruising and bleeding.

Many healthcare providers recommend that patients' TPMT activity levels be tested before thiopurine drugs are prescribed. In people who are found to have reduced enzyme activity, the drugs may be given at a significantly lower dose or different medications can be used to reduce the risk of hematopoietic toxicity.

TPMT deficiency does not appear to cause any health problems other than those associated with thiopurine drug treatment.

Frequency

Studies suggest that less than 1 percent of individuals in the general population have TPMT deficiency. Another 11 percent have moderately reduced levels of TPMT activity that increase their risk of hematopoietic toxicity with thiopurine drug treatment.

Genetic Changes

TPMT deficiency results from changes in the *TPMT* gene. This gene provides instructions for making the TPMT enzyme, which plays a critical role in breaking down

(metabolizing) thiopurine drugs. Once inside the body, these drugs are converted to toxic compounds that kill immune system cells in the bone marrow. The TPMT enzyme "turns off" thiopurine drugs by breaking them down into inactive, nontoxic compounds. Changes in the *TPMT* gene reduce the stability and activity of the TPMT enzyme. Without enough of this enzyme, the drugs cannot be "turned off," so they stay in the body longer and continue to destroy cells unchecked. The resulting damage to the bone marrow leads to potentially life-threatening myelosuppression.

Inheritance Pattern

The activity of the TPMT enzyme is inherited in a pattern described as autosomal codominant. Codominance means that two different versions of the gene are active (expressed), and both versions influence the genetic trait.

The *TPMT* gene can be classified as either low-activity or high-activity. When the gene is altered in a way that impairs the activity of the TPMT enzyme, it is described as low-activity. When the gene is unaltered and TPMT activity is normal, it is described as high-activity. Because two copies of the gene are present in each cell, each person can have two low-activity copies, one low-activity copy and one high-activity copy, or two high-activity copies.

People with two low-activity copies of the *TPMT* gene in each cell have TPMT deficiency and are at the greatest risk of developing hematopoietic toxicity when treated with thiopurine drugs unless they are given much less than the usual dose. People with one high-activity copy and one low-activity copy have moderately reduced enzyme activity and are also at increased risk of this complication unless given a significantly lower dose of the drug. People with two high-activity copies have normal TPMT activity and do not have an increased risk of hematopoietic toxicity with thiopurine drug treatment.

Other Names for This Condition

- poor metabolism of thiopurines
- thiopurine methyltransferase deficiency
- TPMT deficiency

Diagnosis & Management

Genetic Testing

 Genetic Testing Registry: Thiopurine methyltransferase deficiency https://www.ncbi.nlm.nih.gov/gtr/conditions/C0342801/

Other Diagnosis and Management Resources

- MedlinePlus Drug: Azathioprine https://medlineplus.gov/druginfo/meds/a682167.html
- MedlinePlus Drug: Mercaptopurine https://medlineplus.gov/druginfo/meds/a682653.html
- MedlinePlus Drug: Thioguanine https://medlineplus.gov/druginfo/meds/a682099.html

General Information from MedlinePlus

- Diagnostic Tests
 https://medlineplus.gov/diagnostictests.html
- Drug Therapy https://medlineplus.gov/drugtherapy.html
- Genetic Counseling https://medlineplus.gov/geneticcounseling.html
- Palliative Care https://medlineplus.gov/palliativecare.html
- Surgery and Rehabilitation https://medlineplus.gov/surgeryandrehabilitation.html

Additional Information & Resources

MedlinePlus

- Drug: Azathioprine https://medlineplus.gov/druginfo/meds/a682167.html
- Drug: Mercaptopurine https://medlineplus.gov/druginfo/meds/a682653.html
- Drug: Thioguanine https://medlineplus.gov/druginfo/meds/a682099.html
- Encyclopedia: Anemia https://medlineplus.gov/ency/article/000560.htm
- Encyclopedia: Low White Blood Cell Count and Cancer https://medlineplus.gov/ency/patientinstructions/000675.htm
- Encyclopedia: Thrombocytopenia Drug-Induced https://medlineplus.gov/ency/article/000556.htm

- Health Topic: Bone Marrow Diseases https://medlineplus.gov/bonemarrowdiseases.html
- Health Topic: Drug Reactions https://medlineplus.gov/drugreactions.html

Genetic and Rare Diseases Information Center

 Thiopurine S methyltranferase deficiency https://rarediseases.info.nih.gov/diseases/5173/thiopurine-s-methyltranferasedeficiency

Educational Resources

- Leukemia and Lymphoma Society: Normal Blood and Marrow http://www.lls.org/managing-your-cancer/understanding-blood-marrow-and-the-lymphatic-system/normal-blood-and-marrow
- My46 Trait Profile: Thiopurine Response https://www.my46.org/trait-document?trait=Thiopurine%20response&par ent=Medication%20Response&type=profile
- Orphanet: Thiopurine S-methyltransferase deficiency http://www.orpha.net/consor/cgi-bin/OC_Exp.php?Lng=EN&Expert=3315
- St. Jude's Children's Research Hospital: Thiopurine Methyltransferase (TPMT)
 https://www.stjude.org/content/dam/en_US/shared/www/patient-support/do-you-knows/pharmaco-tpmt.pdf

Patient Support and Advocacy Resources

- American Autoimmune Related Diseases Association https://www.aarda.org/
- Leukemia and Lymphoma Society http://www.lls.org/
- Transplant Living from the United Network for Organ Sharing https://transplantliving.org/after-the-transplant/medications/

ClinicalTrials.gov

ClinicalTrials.gov
 https://clinicaltrials.gov/ct2/results?cond=%22thiopurine+S-methyltransferase
 +deficiency%22+OR+%22thiopurine%22

Scientific Articles on PubMed

PubMed

https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28thiopurine+S-methyltran sferase+deficiency%5BTIAB%5D%29+OR+%28TPMT+deficiency%5BTIAB%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+3600+days%22%5Bdp%5D

OMIM

 THIOPURINES, POOR METABOLISM OF, 1 http://omim.org/entry/610460

Sources for This Summary

- Black AJ, McLeod HL, Capell HA, Powrie RH, Matowe LK, Pritchard SC, Collie-Duguid ES, Reid DM. Thiopurine methyltransferase genotype predicts therapy-limiting severe toxicity from azathioprine. Ann Intern Med. 1998 Nov 1;129(9):716-8.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/9841604
- Cooper SC, Ford LT, Berg JD, Lewis MJ. Ethnic variation of thiopurine S-methyltransferase activity: a large, prospective population study. Pharmacogenomics. 2008 Mar;9(3):303-9. doi: 10.2217/14622416.9.3.303.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/18303966
- Coulthard S, Hogarth L. The thiopurines: an update. Invest New Drugs. 2005 Dec;23(6):523-32.
 Review.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/16267626
- Evans WE, Hon YY, Bomgaars L, Coutre S, Holdsworth M, Janco R, Kalwinsky D, Keller F, Khatib Z, Margolin J, Murray J, Quinn J, Ravindranath Y, Ritchey K, Roberts W, Rogers ZR, Schiff D, Steuber C, Tucci F, Kornegay N, Krynetski EY, Relling MV. Preponderance of thiopurine S-methyltransferase deficiency and heterozygosity among patients intolerant to mercaptopurine or azathioprine. J Clin Oncol. 2001 Apr 15;19(8):2293-301.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/11304783
- Fotoohi AK, Coulthard SA, Albertioni F. Thiopurines: factors influencing toxicity and response.
 Biochem Pharmacol. 2010 May 1;79(9):1211-20. doi: 10.1016/j.bcp.2010.01.006. Epub 2010 Jan 21.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/20096268
- Lennard L. Implementation of TPMT testing. Br J Clin Pharmacol. 2014 Apr;77(4):704-14. doi: 10.1111/bcp.12226.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/23962279
 Free article on PubMed Central: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3971986/
- Relling MV, Gardner EE, Sandborn WJ, Schmiegelow K, Pui CH, Yee SW, Stein CM, Carrillo M, Evans WE, Klein TE; Clinical Pharmacogenetics Implementation Consortium. Clinical Pharmacogenetics Implementation Consortium guidelines for thiopurine methyltransferase genotype and thiopurine dosing. Clin Pharmacol Ther. 2011 Mar;89(3):387-91. doi: 10.1038/clpt.2010.320. Epub 2011 Jan 26. Erratum in: Clin Pharmacol Ther. 2011 Dec;90(6):894. Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/21270794
 Free article on PubMed Central: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3098761/

- Schwab M, Schäffeler E, Marx C, Fischer C, Lang T, Behrens C, Gregor M, Eichelbaum M, Zanger UM, Kaskas BA. Azathioprine therapy and adverse drug reactions in patients with inflammatory bowel disease: impact of thiopurine S-methyltransferase polymorphism. Pharmacogenetics. 2002 Aug;12(6):429-36.
 - Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/12172211
- Stanulla M, Schaeffeler E, Flohr T, Cario G, Schrauder A, Zimmermann M, Welte K, Ludwig WD, Bartram CR, Zanger UM, Eichelbaum M, Schrappe M, Schwab M. Thiopurine methyltransferase (TPMT) genotype and early treatment response to mercaptopurine in childhood acute lymphoblastic leukemia. JAMA. 2005 Mar 23;293(12):1485-9.

Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/15784872

Reprinted from Genetics Home Reference:

https://ghr.nlm.nih.gov/condition/thiopurine-s-methyltransferase-deficiency

Reviewed: April 2015

Published: March 21, 2017

Lister Hill National Center for Biomedical Communications U.S. National Library of Medicine National Institutes of Health Department of Health & Human Services